

University of North Georgia
Sophomore Level Mathematics Tournament
April 5, 2014

Solutions for the Afternoon Team Competition

Round 1

Volume = r^2h $6^2 \cdot 3$ $6 \cdot 6 \cdot 3$ $6 \cdot 2 \cdot 3 \cdot 3$ $12 \cdot 9$
The answer is 12 pieces.

Round 2

We think about the complement — people choose different numbers.
The first person can choose any number (positive integer less than 11: from 1 to 10), then the second person would have 9 (different) numbers to choose (9/10), the third person 8 (different) numbers to choose, etc. So the probability that the 4 people choose different numbers is:

$$1 - \frac{9}{10} \cdot \frac{8}{10} \cdot \frac{7}{10} = \frac{504}{1000}$$

Hence the probability that two of the people choose the same number is:

$$1 - \frac{504}{1000} = \frac{496}{1000} = 0.496.$$

Round 3


Since $f(x)$ is divisible by $(x-1)^3$, $x^4 - ax^2 - bx - c = (x-1)^3(x-d)$ for some real number d .

Now if we equate the coefficient of x^3 on both sides we see that $d = 3$.

Then $f(x) = x^4 - 2x^3 + 2x^2 - 3x + 5$.

If you need this document in another format, please email minsu.kim@ung.edu or call 678-717-3546.

Round 4



We get 16 and 8 from the fact that the triangles are congruent. Then we use the Pythagorean Theorem twice getting $a = \sqrt{220} = 2\sqrt{55}$ and $b = \sqrt{55}$. So $a + b = 3\sqrt{55}$.

Round 5

We have $\cot \theta = \cot \phi = 4$, so $\frac{1}{\tan \theta} = \frac{1}{\tan \phi} = 4$ and $\frac{\tan \theta}{\tan \phi} = 4$.

Thus, $\tan \theta = \tan \phi \cdot \frac{\tan \theta}{\tan \phi} = \frac{7}{4}$.

Then

Round 6

Let r be the radius in inches. Then the area in square inches is r^2 which must be a natural number according to the problem.

Since 1.89 T1 0 0 1 93.384 189.05 Tmg.(be)428>DC B1 0 0 1 252.0.1.2/614 gs Es4285ET9.05614 gs EM(oo sq)

Round 7

$f = 1^2 + 2^2 + 4^2 + 6^2 + \dots + 100^2$ and $g = 1^2 + 2^2 + 4^2 + 6^2 + \dots + 98^2 + 100^2$

$g = 1^2 + 3^2 + 5^2 + \dots + 99^2 + 1^2 + 2^2 + 4^2 + 6^2 + \dots + 98^2 + 1^2 + 50^2 + 2^2 + 4^2 + 6^2 + \dots + 98^2$

The sum $2^2 + 4^2 + 6^2 + \dots + 98^2$ can be evaluated as $2^2(1^2 + 2^2 + 3^2 + \dots + 49^2) = 2^2 \cdot 2450 = 2450$.

Consequently, $f = 1^2 + 2450 + 100^2 = 2550$ and $g = 1^2 + 50^2 + 2450 = 2500$.

So $f^2 = 2550^2$ and $g^2 = 2500^2$. $f^2 - g^2 = (f + g)(f - g) = (2550 + 2500)(2550 - 2500) = 252,500$

Dividing 252,500 by 100 gives 2525.

Round 8

We are looking for $abcd = 1200$, where $a, b, c,$ and d are primes with $a < b < c < d$. We solve this problem by finding the largest possible value for a , then for b , and so on. It turns out you can find the answer by making a dozen or so calculations.

- | | |
|---|--------------------------------------|
| | $2 \cdot 3 \cdot 5 \cdot 7 = 210$ |
| 1. Establish a benchmark by multiplying consecutive primes: | $3 \cdot 5 \cdot 7 \cdot 11 = 1155$ |
| | $5 \cdot 7 \cdot 11 \cdot 13 = 5005$ |
- which is the smallest value of $abcd$ where $a = 3$

Round 9

Note that paths cannot be repeated. We will count all the possible paths from S to F that pass through M or N separately and then subtract any paths that are repeated. This is known as an inclusion-exclusion method.

Part 1: Paths from S to F through M (or simply SMF paths) these go from S to M and then to F . There are exactly 3 paths from S to M (of length 3 each). There are exactly 10 paths from M to F (of length 5 each). For each of the 3 SM paths, there are 10 MF paths giving a total of $3 \cdot 10 = 30$ SMF paths.

Part 2: Paths from S to F through N (or simply SNF paths) these go from S to N and then to F . There are 15 paths from S to N (of length 6 each). There are 2 paths from N to F (of length 2 each). For each of the 15 SN paths, there are 2 NF paths giving a tot